

Note

Cd(II)–NCS/NCO complexes of 1-alkyl-2-(arylo)imidazole: Single crystal X-ray structure of [Cd(HaaiMe)₂(SCN)₂]

Kamal Krishna Sarker ^a, Brojo Gopal Chand ^a, Atish Dipankar Jana ^b,
Golam Mostafa ^b, Chittaranjan Sinha ^{a,*}

^a Department of Chemistry, Jadavpur University, Jadavpur, Raja Suboth Mullick Road, Kolkata, West Bengal 700 032, India

^b Department of Physics, Jadavpur University, Jadavpur, Raja Suboth Mullick Road, Kolkata, West Bengal 700 032, India

Received 7 June 2005; received in revised form 1 October 2005; accepted 9 October 2005

Available online 29 November 2005

Abstract

The reaction of Cd(OAc)₂ · 4H₂O and 1-alkyl-2-(arylo)imidazole [RaaiR' where R = H (**a**), Me (**b**); R' = Me (**1/3/5**), Et (**2/4/6**)] and NH₄NCS/NaNCO in methanol in 1:2:2 mole ratio has afforded [Cd(RaaiR')₂(NCS)₂] (**3,4**) and [Cd(RaaiR')₂(NCO)₂] (**5,6**) complexes. The complexes are characterized by different physicochemical methods and in one case, the structure was confirmed by single crystal X-ray diffraction study for title compounds.

© 2005 Elsevier B.V. All rights reserved.

Keywords: Aryloimidazoles; Cadmium(II) complexes; Octahedral; X-ray

1. Introduction

We are interested to design imidazole-containing ligands. There are two basic reasons to concentrate on imidazole type systems: biochemical ubiquity and an alternative to polypyridines. The metal complexes of polypyridines have received much attention because of rich electrochemistry, interesting optical properties, bioinorganic chemistry and catalyses [1–7]. This has led to the modification of the ligand system inserting different substituents, presence of other donor centers etc [7–13]. Ligand synthesis using imidazole as heterocyclic backbone is of much advantage because of the biochemical importance of this molecule [14,15]. Imidazole carries *meta*-related two N centers of different basicity and can bind strongly heavy metal ions [16,17]. We have designed

a surface modified extractant by anchoring imidazole to polystyrene by azo group (–N=N–) and have been used for the separation of heavy metals from drinking water, environmental samples, medicinal samples and ores, minerals [18]. The fundamental property of interaction between metal ions and organic ligands has been utilized in the design of functional materials. Azoimidazoles is one such system. It has also been used to extract anions through protonation of imidazole motif and subsequent interaction with anions via hydrogen bonding [19]. The exo-bidentate behaviour of the ligand has been eliminated by N(1)-alkylation. The coordination chemistry of 1-alkyl-2-(arylo)imidazoles (RaaiR') has been studied by us [20–26] and others [27,28] using transition metals.

In RaaiR', the active function is the azoimine group (–N=N–C=N–) which is isoelectronic with diimine function (–N=C–C=N–). The non-transition metal complexes of aryloimidazole has been reported recently by us [29–32]. This ligand acts as a potential bidentate chelator with

* Corresponding author. Tel.: +91 033 2414 6666x2453; fax: +91 033 2414 6584.

E-mail address: c_r_sinha@yahoo.com (C. Sinha).

heavier transition metals [20–26]. It comprises with uni- or bidenticity when coordinated with 3d block [20–22] and non-transition metals [29–32]. The Group 12 (Zn, Cd, Hg) metal complexes do not have CFSE and the stereochemistry of the complexes has been controlled by steric effect and the M–N bond strength. As a part of our programme to explore the coordination chemistry of Group 12 metal complexes of arylazoimidazoles, we wish to report in this work the Cd(II)-thiocyanato/isocyanato complexes of 1-alkyl-2-(arylo)imidazoles (RaaiR'). The X-ray structure of one of the complexes has been described.

2. Experimental

2.1. Materials

Published methods [20] were used to prepare 1-alkyl-2-(arylo)imidazoles (RaaiR' where R = H (**a**), Me (**b**); R' = Me (**1,3,5**), Et (**2,4,6**)). All other chemicals and organic solvents used for preparation work were of reagent grade received from SRL, India.

2.2. Physical measurements

Microanalyses (C, H, N) were performed using a Perkin–Elmer 2400 CHNO/S elemental analyzer. Spectroscopic measurements were carried out using the following instruments: UV–Vis spectra, reflectance spectra, JASCO UV–Vis/NIR model V-570; IR spectra (KBr disk, 4000–200 cm⁻¹), JASCO FT-IR model 420.

2.3. Preparation of bis-thiocyanato bis-[1-methyl-2-(phenylazo)imidazole] cadmium(II)

2.3.1. [Cd(HaaiMe)₂(NCS)₂] (**3a**)

Methanolic solution (15 ml) of 1-methyl-2-(phenylazo)imidazole (HaaiMe, 0.22 g, 1.18 mmol) was added dropwise to Cd(OAc)₂·4H₂O (0.15 g, 0.5 mmol) in the same solvent (20 ml) at room temperature (298 K). Then aqueous-methanolic solution of NH₄NCS (0.08 g, 1.05 mmol) was added dropwise. The mixture was stirred for 15 min and filtered. Filtrate was kept undisturbed for few days. Orange-red crystals were obtained and washed with cold water and methanol. Finally, they were dried in vacuo. Isolated yield was 0.185 g, (62%).

All other complexes were prepared by the same procedure and yield varied 60–70%. Microanalytical data of the complexes are as follows: [Cd(HaaiMe)₂(NCS)₂] (**3a**): *Anal. Calc.* for C₂₂H₂₀N₁₀S₂Cd: C, 43.96; H, 3.33; N, 23.31. Found: C, 43.90; H, 3.27; N, 23.38%. [Cd(MeaaiMe)₂(NCS)₂] (**3b**): *Anal. Calc.* for C₂₄H₂₄N₁₀S₂Cd: C, 45.82; H, 3.81; N, 22.27. Found: C, 45.73; H, 3.74; N, 22.18%. [Cd(HaaiEt)₂(NCS)₂] (**4a**): *Anal. Calc.* for C₂₄H₂₄N₁₀S₂Cd: C, 45.82; H, 3.81; N, 22.27. Found: C, 45.76; H, 3.74; N, 22.18%. [Cd(MeaaiEt)₂(NCS)₂] (**4b**): *Anal. Calc.* for

C₂₆H₂₈N₁₀S₂Cd: C, 47.53; H, 4.26; N, 21.32. Found: C, 47.44; H, 4.18; N, 21.24%.

2.4. Preparation of bis-isocyanato-bis-[1-methyl-2-(phenylazo)imidazole] cadmium(II)

2.4.1. [Cd(HaaiMe)₂(NCO)₂] (**5a**)

Methanolic solution (15 ml) of 1-methyl-2-(phenylazo)imidazole (HaaiMe, 0.22 g, 1.18 mmol) was added dropwise to Cd(OAc)₂·4H₂O (0.15 g, 0.5 mmol) in the same solvent (20 ml) at room temperature (298 K). Then aqueous-methanolic solution of NaNCO (0.07 g, 1.08 mmol) was added dropwise. The mixture was stirred for 15 min and filtered. Filtrate was kept undisturbed for few days. Orange-red crystals were obtained and washed with cold water and methanol. Finally they were dried in vacuo. Isolated yield was 0.15 g, (53%).

All other complexes were prepared by the same procedure and yield varied 60–70%. Microanalytical data of the complexes are as follows: [Cd(HaaiMe)₂(NCO)₂] (**5a**): *Anal. Calc.* for C₂₂H₂₀N₁₀O₂Cd: C, 46.45; H, 3.52; N, 24.63. Found: C, 46.32; H, 3.60; N, 24.48%. [Cd(MeaaiMe)₂(NCO)₂] (**5b**): *Anal. Calc.* for C₂₄H₂₄N₁₀O₂Cd: C, 48.29; H, 4.02; N, 23.47. Found: C, 48.18; H, 3.94; N, 23.33%. [Cd(HaaiEt)₂(NCO)₂] (**6a**): *Anal. Calc.* for C₂₄H₂₄N₁₀O₂Cd: C, 48.29; H, 4.02; N, 23.47. Found: C, 48.20; H, 3.90; N, 23.35%. [Cd(MeaaiEt)₂(NCO)₂] (**6b**): *Anal. Calc.* for C₂₆H₂₈N₁₀O₂Cd: C, 49.97; H, 4.48; N, 22.42. Found: C, 49.81; H, 4.43; N, 22.35%.

2.5. X-ray crystal structure analysis of [Cd(HaaiMe)₂(NCS)₂] (**3a**)

Crystal is growing (orange block, 0.3 × 0.3 × 0.2 mm³) by slow evaporation of methanolic solution of mixture containing Cd(OAc)₂, HaaiMe and NH₄CNS in 1:2:2 molar ratio. A summary of the crystallographic data and structure refinement parameters are given in Table 1. Data were collected with Siemens SMART CCD diffractometer using graphite-monochromatized Mo K α radiation ($\lambda = 0.71073$ Å) at 293(2) K. The intensity data were corrected for Lorentz and polarisation effects and an empirical absorption correction was employed using SAINT program. A total of 16633 reflections [range 3.4 ≤ 2 θ ≤ 56.74°] were collected and 3600 were assumed observed applying the condition $I > 2\sigma(I)$. The structure was solved by direct methods and followed by successive Fourier and difference Fourier syntheses. Full matrix least squares refinements on F^2 were carried out for both cases using SHELXL-97 with anisotropic displacement parameters for all non-hydrogen atoms. The hydrogen atoms were fixed geometrically and refined using the riding model. In the final difference Fourier map the residual minima and maxima were –0.554 and 0.939 e/Å³. Complex neutral atom scattering factors were used throughout. All calculations were carried out using SHELXS-97, SHELXL-97, PLATON-99, ORTEP programs.

Table 1
Summarized crystallographic data for [Cd(HaaiMe)₂(NCS)₂] (**3a**)

[Cd(HaaiMe) ₂ (NCS) ₂] (3a)	
Empirical formula	C ₂₂ H ₂₀ CdN ₁₀ S ₂
Formula weight	601.03
Temperature (K)	293
Crystal system	monoclinic
Space group	P2 ₁ /c (No. 14)
<i>Unit cell dimensions</i>	
<i>a</i> (Å)	8.9960(9)
<i>b</i> (Å)	15.3217(15)
<i>c</i> (Å)	19.352(2)
β (°)	91.445(2)
<i>V</i> (Å ³)	2666.5(5)
<i>Z</i>	4
λ (Å)	0.71073
μ (Mo K α) (mm ⁻¹)	1.006
<i>D</i> _{calc} (mg m ⁻³)	1.497
Refine parameters	316
Total collected data	16633
Unique data [<i>I</i> > 2 σ (<i>I</i>)]	3600
<i>R</i> ₁ ^a [<i>I</i> > 2 σ (<i>I</i>)]	0.0760
<i>wR</i> ₂ ^b	0.2644
Goodness-of-fit	1.03

$$^a R = \sum |F_o - F_c| / \sum F_o$$

$$^b wR = [\sum w(F_o^2 - F_c^2) / \sum wF_o^4]^{1/2} \text{ are general but } w = 1 / [\sigma^2(F_o^2) + (0.1616P)^2 + (0.0000P)], \text{ where } P = (F_o^2 + 2F_c^2) / 3.$$

3. Result and discussion

3.1. Synthesis and formulation

The reaction of Cd(OAc)₂ · 4H₂O, RaaiR' and NH₄CNS in the mole proportion of 1:1:2 has synthesized $\mu_{1,3}$ -NCS bridged cadmium polymer [30]. The reaction of Cd(ClO₄)₂ or Cd(NO₃)₂ salt with excess of RaaiR', however synthesized tetrahedral Cd(RaaiR')₄²⁺ compounds [32]. In this work we are using different compositions of the metal:ligand:NCS⁻. Methanolic solution of Cd(OAc)₂ · 4H₂O and RaaiR' in the mole ratio of 1:2 followed by the addition of 2 equivalent of NH₄NCS or NaNCO at later stage has isolated complexes of the composition [Cd(RaaiR')₂(NCS)₂] (**3, 4**) and [Cd-(RaaiR')₂(NCO)₂] (**5, 6**), respectively. The microanalytical data support the composition of the complexes. They are non-electrolyte in nature.

3.2. Spectral studies

3.2.1. IR spectra

In the infrared spectra, the point of interest is the band due to the thiocyanate (–NCS)/isocyanate (–NCO), azo (–N=N–) and imine (–C=N–) groups in the complexes. The spectra show very strong band at 2063–2067 cm⁻¹ for **3** and **4** which are assigned to ν (NCS) (Table 2). The compounds **5, 6** show very strong band at 2190–2208 cm⁻¹ and have been assigned to ν (NCO). Moderately intense stretchings at ca. 1595 and ca. 1445 cm⁻¹ are due to ν (C=N) and ν (N=N), respectively. In free ligands [22,29]

ν (N=N) and ν (C=N) appear at 1455–1460 and 1605–1615 cm⁻¹. In the complexes, these are shifted to lower frequency region which are in support of coordination of azo-N and imine-N to Cd(II) centre.

3.2.2. Electronic spectra

The solution electronic spectra of the complexes **3** and **4** show intense transition <400 nm and are assigned to ligand centered transitions. Two weak ($\epsilon \sim 10^3$ dm³ mol⁻¹ cm⁻¹) transitions are observed at >440 nm (Table 2). The ligands are π -acidic and Cd²⁺ is a d¹⁰ system, thus the transition may be assigned to d(Cd) → π^* (ligand) transition.

3.2.3. ¹H NMR spectra

The ¹H NMR spectra of [Cd(RaaiR')₂(NCS)₂] (**3, 4**) and [Cd(RaaiR')₂(NCO)₂] (**5, 6**) were recorded in CD₃CN. The spectral data are collected in Table 3. The proton numbering pattern is shown in Scheme 1. The assignment has been made on the basis of spin–spin interaction and on comparing with reported data [30–32]. Imidazole protons (4- and 5-H) of [Cd(RaaiR')₂(NCS)₂] (**3, 4**) appear at ca. 7.5 and 7.3 ppm and are singlet in nature which may be due to rapid proton exchange with solvent proton (CD₃CN is hydrolysed on exposure to air). In Cd(RaaiR')₂(NCO)₂ (**5, 6**) 4-H appears at ~0.3 ppm (ca. 7.6 ppm) higher δ than 5-H (ca. 7.3 ppm). It is concluded that 4-H is closer to the metal centre. This is also supported from single crystal X-ray structure (Fig. 2) where RaaiR' is bounded to Cd(II) through N(3) center only. Influence of anion ligand is also reflected from the NMR data; being a better electron withdrawing agent NCO than NCS, the chemical shift of the coordinated ligands in **5, 6** are shifted to higher δ by about 0.1 ppm. N(1)–Me appears at ca. 4 ppm; N(1)–CH₂–CH₃ shows quartet and triplet splitting signals at ca. 4.4 (*J* = 9.0 Hz) and ca. 1.6 (*J* = 7.5 Hz) ppm, respectively, for –CH₂, and –CH₃ groups. The N(1)–CH₂–Ph gives a singlet response at ~5.35 ppm. Aryl protons (7-11-H) appear at usual position (7.4–7.8 ppm) and suffer shifting to down-field position with reference to free ligand value.

3.3. Molecular structure

The molecular structure of [Cd(NCS)₂(HaaiMe)₂] (**3a**) is shown in Fig. 1. Bond distances and angles are set out in Table 4. The metric parameters (Table 2) and figure show distorted CdN₆ coordination sphere. The octahedral coordination is satisfied by four N centers from two HaaiMe ligands and other two N-centres from two NCS groups. HaaiMe acts as bidentate N(imidazole), N(azo) chelator. The chelating N–Cd–N bond angle is N(1)–Cd(1)–N(4), 67.2(2)° and lies in the limiting value of Group 12 complexes of azoheterocycles [29–32]. Other chelate angle N(5)–Cd(1)–N(8) is very low, 62.2(1)°. Very long Cd(1)–N(8) bond distance (2.919(2) Å) may be due to the low chelate angle. In pseudo-octahedral compounds of type MX₂(N,N')₂, five different isomers are possible: two compounds having *trans*-MX₂ configuration and three

Table 2
UV–Vis spectra^a, and IR^b

Compound	UV–Vis spectral data (λ_{\max}/nm) ($10^{-3} \epsilon \text{ M}^{-1} \text{ cm}^{-1}$)	IR spectra, cm^{-1}		
		$\nu(\text{NCS}/\text{NCO})$	$\nu(\text{C}=\text{N})$	$\nu(\text{N}=\text{N})$
[Cd(HaaiMe) ₂ (NCS) ₂] (3a)	432(0.99), 386(2.81), 372(2.77), 232(1.33)	2060	1590	1440
[Cd(MeaaiMe) ₂ (NCS) ₂] (3b)	442(1.69), 384(3.74), 372(2.83), 234(1.22)	2060	1598	1444
[Cd(HaaiEt) ₂ (NCS) ₂] (4a)	432(1.42), 390(3.45), 356(3.97), 236(1.67)	2055	1585	1445
[Cd(MeaaiEt) ₂ (NCS) ₂] (4b)	434(1.51), 386(3.29), 370(3.20), 244(1.09)	2060	1595	1440
[Cd(HaaiMe) ₂ (NCO) ₂] (5a)	445(0.73), 395(2.65), 370(2.67), 238(1.40)	2190	1600	1448
[Cd(MeaaiMe) ₂ (NCO) ₂] (5b)	452(0.69), 398(3.45), 370(2.73), 236(1.32)	2202	1594	1450
[Cd(HaaiEt) ₂ (NCO) ₂] (6a)	438(1.28), 396(3.55), 350(3.75), 240(1.74)	2208	1605	1455
[Cd(MeaaiEt) ₂ (NCO) ₂] (6b)	444(1.21), 392(3.16), 360(3.10), 244(1.26)	2195	1600	1450

^a In CH₃OH.

^b In KBr disk.

Table 3
¹H NMR spectral data in CD₃CN at room temperature


Compound	δ , ppm (<i>J</i> , Hz)							
	4-H ^a	5-Ha	7,11-Ha	8-10-H	9-Me	1-Me	1-CH ₂ ^c	(1-CH ₂)CH ₃ ^d
3a	7.48(7.0)	7.27(7.0)	7.90(7.5)	7.44 ^b		4.09		
3b	7.50(7.0)	7.27(7.0)	7.92(7.5)	7.34(7.5) ^a	2.45	4.11		
4a	7.53(7.5)	7.28(7.5)	7.91(7.8)	7.46 ^b			4.42 (12.0)	1.52 (7.5)
4b	7.51(7.5)	7.32(7.5)	7.89(7.5)	7.34(7.5) ^a	2.45		4.50 (12.0)	1.57 (7.5)
5a	7.61(7.0)	7.30(7.0)	7.92(7.5)	7.55 ^b		4.10		
5b	7.64(7.5)	7.32(7.5)	7.95(7.8)	7.37(7.0) ^a	2.47	4.13		
6a	7.62(7.0)	7.30(7.0)	7.95(7.0)	7.51 ^b			4.04 (10.0)	1.55 (7.0)
6b	7.59(7.0)	7.29(7.0)	7.92(7.0)	7.34(7.0) ^a			4.52 (10.0)	1.58 (7.0)

^a Doublet.

^b Multiplet.

^c Quartet.

^d Triplet.



Scheme 1. RaaiR': R = H (1,3,5); Me (2, 4, 6) R' = Me (a), Et (b).

compounds having *cis*-MX₂ configuration [25]. Sequence of coordination of pairs of donor centers following pairs of X, N and N' donor centers the isomers are assigned as *trans*-*cis*-*cis*, *trans*-*trans*-*trans*, *cis*-*trans*-*trans*, *cis*-*trans*-*cis*, and *cis*-*cis*-*cis*. The orientation of donor centers in the present complex about Cd(II) has constituted *cis*-*cis*-*cis* type geometrical isomer. The atomic arrangements Cd(1), N(1), C(1), N(3), N(4); Cd(1), N(5), C(12), N(7), N(8) constitute two chelate planes (deviation < 0.03 Å) and are inclined at dihedral ca. 70°. The pendant phenyl groups are also planar and make dihedral of ca. 7° with respective chelated azoimine fragment. Three atomic




Fig. 1. ORTEP diagram of [Cd(HaaiMe)₂(NCS)₂] (with 50% probability ellipsoids).

arrangements Cd(1), N(4), N(5), N(8), N(9) (plane-1); Cd(1), N(1), N(8), N(9), N(10) (plane-2); Cd(1), N(4), N(5), N(10) (plane-3) constitute three molecular

Table 4
Selected bond distances and bond angles of **3a**

Bond distance (Å)		Bond angle (°)	
Cd(1)–N(1)	2.086(6)	N(1)–Cd(1)–N(4)	67.2(2)
Cd(1)–N(4)	2.677(5)	N(1)–Cd(1)–N(5)	105.4(2)
Cd(1)–N(5)	2.202(5)	N(1)–Cd(1)–N(9)	111.2(3)
Cd(1)–N(9)	2.124(8)	N(1)–Cd(1)–N(10)	130.9(3)
Cd(1)–N(10)	2.044(6)	N(4)–Cd(1)–N(5)	171.81(17)
S(1)–C(20)	1.616(8)	N(4)–Cd(1)–N(9)	87.8(2)
S(2)–C(19)	1.608(7)	N(4)–Cd(1)–N(10)	86.5(2)
N(9)–C(20)	1.137(11)	N(5)–Cd(1)–N(9)	92.0(2)
N(10)–C(19)	1.120(9)	N(5)–Cd(1)–N(10)	101.3(2)
N(3)–N(4)	1.243(8)	N(9)–Cd(1)–N(10)	108.2(3)
N(7)–N(8)	1.253(7)	Cd(1)–N(4)–N(3)	111.4(4)
Cd(1)–N(8)	2.919(2)	N(8)–Cd(1)–N(5)	62.2(1)
		N(8)–Cd(1)–N(9)	154.1(7)
		N(8)–Cd(1)–N(10)	78.0(2)
		N(8)–Cd(1)–N(1)	79.1(3)
		N(8)–Cd(1)–N(4)	117.9(3)

planes and maximum deviation has been observed for N(azo) (N(4/8)) and they make dihedral 75–88°.

There are three types of N-centres: N(imidazole) (N(1)/N(5)), N(azo) (N(4)/N(8)) and N(thiocyanato) (N(9)/N(10)). The Cd–N(thiocyanato) and Cd–N(imidazole) distances do not differ widely. The Cd–N(azo) distances

are the longest in the series and they are not equivalent: Cd(1)–N(4), 2.677(5) and Cd(1)–N(8), 2.919(2) Å. It is because the N(azo) requires considerably more room than the other two types of N-donors. There are at least two steric reasons for this. First, the N(azo) belongs to exocyclic N=N along with a pendant phenyl group; thus the van der Waals repulsion will be greater than N(imidazole). The lowering in overall symmetry is manifested by the elongation of two Cd–N(azo) distances. However, the Cd–N(azo) distances are less than the sum of the van der Waals radii of Cd(II) (1.58 Å) and N(azo) (1.55 Å) [33,34]. This implies covalent interaction of N(azo) and Cd(II).

Three different types of non-covalent interactions (C–H–S; C–H– π and π – π) are observed in the packing view of the molecule (Figs. 2–4). Hydrogen bonding with non-bonded S is scarce [35]. The coordinated M–NCS is hydrogen bonded with H of N–CH₃ group and 1-D chain has been generated along the *x*-direction [C(22)–H(22b)–S(1): C–H, 0.96 Å; H–S(1), 2.76 Å; C(22)–S(1), 3.602(10) Å; \angle C(22)–H(22b)–S(1), 164°; symmetry: *x*, 1/2 – *y*, 1/2 + *z*] (Fig. 2). It is proposed that S in M–NCS is more electronegative and less polarisable than sulfide-S and may exhibit an affinity to acidic-H. In fact, this proton abstracting affinity in dilute acid solution is




Fig. 2. 1-D chain constituted by hydrogen bonding along *x*-direction.




Fig. 3. Sheet formation of 1D chain by C–H– π interaction.




Fig. 4. Unit cell pack showing π - π interaction between inter-chain pendant phenyl groups.

responsible for liberation of H_2S from thiourea, thiocarbamate [36] and serves as a very good supplement of H_2S gas in analytical laboratory for environmental reasons. The C-H- π interaction is also observed in C(22)-H(22A)-Cg(5): H-Cg, 2.75 Å; \angle C(22)-H(22A)-Cg(5), 164°; where Cg(5) stands for pendant phenyl ring, C(13), C(14), C(15), C(16), C(17), C(18) (symmetry: $1+x, y, z$) (Fig. 3). An inter-chain π - π interaction is observed between pendant Ph rings, Cg(4)-Cg(9), 3.887(4) Å (Cg(4): C(4), C(5), C(6), C(7), C(8), C(9); Cg(9): C(4a), C(5a), C(6a), C(7a), C(8a), C(9a) where a stands for the neighboring molecule symmetry, $1-x, -y, 1-z$) which constitutes a supramolecular structure (Fig. 4).

4. Supplementary material

Crystallographic data for the structural analysis has been deposited with the Cambridge Crystallographic data center, CCDC No. 274228 for $[\text{Cd}(\text{Haa}i\text{Me})_2(\text{NCS})_2]$ (**3a**). Copies of this information may be obtained free of charge from the Director, CCDC, 12 Union Road, Cambridge, CB2 1FZ, UK.

Acknowledgement

Financial supports (C.S.) from the Council of Scientific and Industrial Research, New Delhi are gratefully acknowledged. One of us (U.S.R.) also thanks the UGC for a fellowship. Our sincere thanks are due to Prof. T.-H. Lu, Department of Physics, National Tsing Hua University, Taiwan, ROC for helpful discussion.

References

- [1] J. Reedijk, in: G. Wilkinson, J.A. McCleverty (Eds.), *Comprehensive Coordination Chemistry*, vol. 2, Pergamon Press, Oxford, UK, 1987, p. 43.
- [2] F.A. Cotton, G. Wilkinson, *Advanced Inorganic Chemistry*, fifth ed., Wiley International Science, 1994.
- [3] K.E. Erkkila, D.T. Odam, J.K. Barton, *Chem. Rev.* 99 (1999) 2777.
- [4] C. Kaes, A. Katz, M.W. Hosseini, *Chem. Rev.* 100 (2000) 3553.
- [5] J.M. Lehn, *Supramolecular Chemistry*, VCH, Weinheim, 1995.
- [6] K. Kalyansundaram, *Coord. Chem. Rev.* 46 (1982) 159.
- [7] W. Guo, Z. Peng, D. Li, Y. Zhou, *Polyhedron* 23 (2004) 1701, and references therein.
- [8] Y.-H. Kim, J.S. Youk, S.Y. Moon, J.-I. Choc, S.-K. Chng, *Chem. Lett.* 33 (2004) 702.
- [9] L.G. Nikolchava, C.M. Vogels, R.A. Stefan, H.A. Darwish, S.J. Duffy, R.J. Ireland, A. Decken, R.H.E. Hudson, S.A. Wertcott, *Can. J. Chem.* 81 (2003) 269.
- [10] M. Menon, A. Pramanik, N. Bag, A. Chakravorty, *Inorg. Chem.* 33 (1994) 403.
- [11] E.C. Constable, *Coord. Chem. Rev.* 93 (1989) 205.
- [12] R. Mukherjee, *Coord. Chem. Rev.* 203 (2000) 151.
- [13] B.K. Ghosh, A. Chakravorty, *Coord. Chem. Rev.* 95 (1989) 239.
- [14] H. Sigel (Ed.), *Metal Ions in Biological System*, different vols., Marcel Dekker, New York, 1981.
- [15] A. Butler, C.J. Carrano, *Coord. Chem. Rev.* 109 (1991) 61.
- [16] W. Kaim, B. Schwederski, *Bioinorganic Chemistry: Inorganic Elements in the Chemistry of Life*, Wiley, New York, 1994.
- [17] S.J. Lippard, in: I. Bertini, H.B. Gray, S.J. Lippard, J.C. Valentine (Eds.), *Bioinorganic Chemistry*, University Science Books, Mill Valley, 1994.
- [18] D. Das, A.K. Das, C. Sinha, *Talanta* 48 (1999) 1013.
- [19] U.S. Ray, B.G. Chand, A.K. Dasmahapatra, G. Mostafa, T.-H. Lu, C. Sinha, *Inorg. Chem. Commun.* 6 (2003) 634–638.
- [20] U.S. Ray, B.G. Chand, G. Mostafa, J. Cheng, T.-H. Lu, C. Sinha, *Polyhedron* 22 (2003) 2587.
- [21] U.S. Ray, D. Banerjee, G. Mostafa, T.-H. Lu, C. Sinha, *New. J. Chem.* 28 (2004) 1.
- [22] D. Banerjee, U.S. Ray, J.-C. Liou, C.-N. Lin, T.-H. Lu, C. Sinha, *Inorg. Chim. Acta* 358 (2005) 1019, and references therein.
- [23] Sk. Jasimuddin, P. Byabartta, G. Mostafa, J.-C. Liou, T.-H. Lu, C. Sinha, *J. Coord. Chem.* 57 (2004) 75.
- [24] J. Dinda, P.K. Santra, C. Sinha, L.R. Falvello, *J. Organomet. Chem.* 629 (2001) 28.
- [25] P. Byabartta, S. Pal, C. Sinha, F.-L. Liao, K. Panneerselvam, T.-H. Lu, *J. Coord. Chem.* 55 (2002) 479.
- [26] S. Pal, D. Das, C. Sinha, C.H.L. Kennard, *Inorg. Chim. Acta* 313 (2001) 21.
- [27] M.N. Ackermann, M.P. Robinson, I.A. Maher, E.B. LeBlanc, R.V. Raz, *J. Organomet. Chem.* 682 (2003) 248.
- [28] I. Chakravorty, S. Sengupta, S. Das, S. Banerjee, A. Chakravorty, *J. Chem. Soc., Dalton Trans.* 134 (2003).
- [29] B.G. Chand, U.S. Ray, G. Mostafa, T.-H. Lu, C. Sinha, *J. Coord. Chem.* 57 (2004) 627–634.
- [30] B.G. Chand, U.S. Ray, G. Mostafa, T.-H. Lu, C. Sinha, *Polyhedron* 23 (2004) 1669.
- [31] B.G. Chand, U.S. Ray, P.K. Santra, G. Mostafa, T.-H. Lu, C. Sinha, *Polyhedron* 22 (2003) 1205.
- [32] B.G. Chand, G. Mostafa, T.-H. Lu, L.R. Falvello, C. Sinha, *Polyhedron* 22 (2003) 3161.
- [33] A. Bondi, *J. Phys. Chem.* 68 (1964) 441.
- [34] A.M. James, M.P. Lord, *Macmillan's Chemical and Physical Data*, Macmillan, London, UK, 1992.
- [35] Y.-B. Jiang, H.-Z. Kou, F. Gao, R.-J. Wang, *Acta Crystallogr.* c60 (2004) m261.
- [36] A.I. Vogel, *A Text Book of micro and Semimicro Qualitative Inorganic Analyses*, fourth ed., Orient Longmans Ltd, 1962.